

설기저부의 소타액선에 발생한 투명세포선암종 1례

대전선병원 이비인후과,¹ 고려대학교 의과대학 이비인후-두경부외과학교실²
조승현¹ · 박규선¹ · 김석현¹ · 권순영²

A Case of Clear Cell Adenocarcinoma on Tongue Base

Syung-Hyun Cho, MD¹, Kue-Sun Park, MD¹, Suk-Hyun Kim, MD¹ and Soon-Young Kwon, MD²¹Department of Otolaryngology, Sun General Hospital, Daejeon²Department of Otolaryngology-Head and Neck Surgery, College of Medicine, Korea University, Seoul, Korea

-ABSTRACT-

It is reported that tumors occurring on minor salivary glands represent less than 10 percent of total salivary gland tumors. In particular, the occurrence of clear cell adenocarcinoma is very rarely observed. Thus, it is hard to measure the exact occurrence rate. Nevertheless, it is assumed that clear cell adenocarcinoma covers less than one percent of all salivary glands tumors. Clear cell adenocarcinoma shows similar histological morphology with salivary gland tumors such as mucopidermoid carcinoma, acinic cell carcinoma, epithelial and myoepithelial tumors. However, its differential diagnosis with other tumors is possible because it shows the unique monomorphous clear cells. Authors observed the clear cell adenocarcinoma of minor salivary glands at tongue base in a 60 year-old female patient who was complaining about a foreign body sensation on the pharynx. (J Clinical Otolaryngol 2002;13:251-253)

KEY WORDS : Adenocarcinoma · Clear cell · Salivary glands · Minor.

서론

(monomorphous)¹⁾

60

10%

1%

증례

2000 8 60

가

9

2000 2

가

2 x 2 cm

(Fig. 1),

: 2002 9 5

: 2002 10 11

: , 425 - 020 516

: (031) 412 - 5170 · : (031) 401 - 4365

E - mail : entkwon@chollian.net



Fig. 1. Endoscopic finding of the cancer of tongue base. A protruding mass, about 2 × 2 cm sized, on tongue base is seen.




Fig. 2. CT scan with contrast enhancement. about 3 × 2 cm sized, enhanced soft tissue density mass at right tongue base is demonstrated.



Fig. 3. The surgical specimen. The specimen consists of a portion of the tongue, measuring 4.5 × 4.5 cm with mass, measuring 3 cm in diameter.




Fig. 4. Pathologic finding. Tumor tissue is composed of nests and cords of monomorphous cells with clear cytoplasm that have well defined cell boundaries and eccentrically located nuclei.

cm

(Fig. 2). 2000 9 6

3 × 2 cm

가

가

3 × 2

가

(nest)

(cord)

가

(Fig.

4),

가

24

고 찰

,

,

-

가

가

¹⁾²⁾ 1991 AFIP(Armed Forces Institute of Pathology) Alternative Terminology Used in WHO Classification

가

1%

60%가

40 70

15

¹⁾

가

가

³⁾

⁴⁾⁵⁾ (glycogen) PAS (sheet),

가

¹⁾²⁾¹¹⁾ 가

cytokeratin S-100

GFAP(Glial Fibrillary Acidic Protein), actin, vimentin

¹⁾⁴⁾

가 ¹⁾¹¹⁾

중심 단어 :

REFERENCES

- 1) Auclair PL, Ellis GL. *Major Salivary Glands*: In Silverberg SG, Delellis RA, Frable WJ, editor. *Principles and Practice of Surgical Pathology and Cytopathology*. 3rd ed. Churchill Livingstone Inc. 1997;33:1462-91.
- 2) Chen KT. *Clear cell carcinoma of the salivary gland*. *Hum Pathol* 1983;14:91-3.
- 3) Milchgrub S, Gnepp DR, Vuitch F, Dolgado R, Albores-Saavedra J. *Hyalinizing clear cell carcinoma of salivary gland*. *Am J surg Pathol* 1994;18:74-82.
- 4) Ogawa I, Nikai H, Takata T, Ijuhin N, Miyauchi M, Ito H, et al. *Clear cell tumors of minor salivary gland origin. An immunohistochemical and ultrastructural analysis*. *Oral Surg Oral Med Oral Pathol* 1991;72:200-7.
- 5) Simpson RH, Sarsfield PT, Clarke T, Babajews AV. *Clear cell carcinoma of minor salivary glands*. *Histopathology* 1990;17:433-8.
- 6) Lattanzi DA, Polverini P, Chin DC. *Glycogen-rich adenocarcinoma of a minor salivary gland*. *J Oral Maxillofac Surg* 1985;43:122-4.
- 7) Chaudhry AP, Cutler LS, Satchidanand S, Labay G, Raj MS, Lin CC. *Glycogen-rich tumor of the oral minor salivary glands. A histochemical and ultrastructural study*. *Cancer* 1983;52:105-11.
- 8) Hayashi K, Ohtsuki Y, Sonobe H, Takahashi K, Iwata J, Nishioka E, et al. *Glycogen-rich clear cell carcinoma arising from minor salivary glands of the uvula. A case report*. *Acta Pathol Jpn* 1988;38:1227-34.
- 9) Uri AK, Wetmore RF, Iozzo RV. *Glycogen-rich clear cell carcinoma in the tongue. A cytochemical and ultrastructural study*. *Cancer* 1986;57:1803-9.
- 10) Mohamed AH, Cherrick HM. *Glycogen-rich adenocarcinoma of minor salivary glands. A light and electron microscopic study*. *Cancer* 1975;36:1057-66.
- 11) Eversole LR. *On the differential diagnosis of clear cell tumors of the head and neck*. *Eur J Cancer B Oral Oncol* 1993; 29B:173-9.