

## 일측성 말초 전정계를 침범한 신경베체트병 치험 1예

한기훈 · 김위황 · 김남걸 · 손원상

A Case of Unilateral Peripheral Vestibular Involvement  
in Neuro-Behçet's Disease

Ki Hun Han, MD, Wee Hwang Kim, MD, Nam Gol Kim, MD and Won Sang Son, MD

Department of Otolaryngology, Gwangju Christian Hospital, Gwangju, Korea

## -ABSTRACT -

Behçet's disease is a chronic, multisystemic disorder that affects mainly young adults and is characterized by a relapsing inflammatory process of unknown cause. It was originally described as a clinical triad of oral aphthae, genital ulcers, ocular lesions. Furthermore, cutaneous, central nervous system, vascular and cardiovascular, gastrointestinal, renal, and pleuropulmonary lesions, arthritis, and epididymitis are today accepted as additional clinical features of Behçet's syndrome. Hearing loss and dizziness are frequent symptoms in patients affected by Behçet's syndrome with or without the involvement of the CNS. Peripheral involvement is more common than central involvement for vestibulocochlear lesions. In addition, vestibular deficit may represent an early sign of neuro-Behçet's syndrome (CNS lesion in behçet's disease). We experienced unilateral peripheral vestibular involvement without cochlear involvement in neuro-Behçet's syndrome, so we report this case with a review of literature. (J Clinical Otolaryngol 2005;16:140-143)

KEY WORDS : Behçet's syndrome · Vertigo.

## 서 론

62~80%

3)5)

10~49%

7)

3가

가

8)

1-6)

3)9)10)

: 2005 4 6

: 2005 5 17

: , 503 - 715

264

## 증 례

: (062) 650 - 5095 · : (062) 650 - 5090

E - mail : kchent2@hanmail.net

36

1

2 ANA, VDRL

Pathergy HLA - B51

2

4

Frenzel

Romberg Mann

(Fig. 1).


3

4

Hamid 5


Pathergy

HLA - B51



**Fig. 1.** Electronystagmography shows spontaneous nystagmus to left. Caloric test shows evidence of right side canal paresis.

**Fig. 2.** A : Contrast-enhanced axial T1-weighted MR image : Meningeal enhancement (arrowhead) and transverse sinus thrombosis (arrow) was seen. B : Contrast-enhanced T1-weighted coronal image : Leptomeningeal enhancement (arrowhead) and sigmoid sinus thrombosis (arrow).



가 Gemignani <sup>6)</sup> 16  
 S 8 , 3  
 (leptomeninges) , 5  
 (Fig. 2),  
 3 solumedrol . Elidan <sup>5)</sup>  
 1.0 g pulse therapy cyclosporine 750 mg 35 28 (80%)  
 prednisolon 50 mg 8 , 20  
 10 가 Pollak <sup>12)</sup> 26  
 10 , 19 (73%)  
 , 15  
 6

**고 찰**

10~49% <sup>7)</sup>  
 가  
<sup>4)11)</sup>  
 가 10 <sup>3)9)11)</sup>  
 가  
 S <sup>12)</sup>  
<sup>6)</sup>  
 1961 Ala- 가  
 jouanine <sup>10)</sup>  
 가  
 Brama <sup>3)</sup> 16 10 , , ,  
 6 , , , ,  
 가 <sup>2)</sup>

중심 단어 :

가 가 1)3)7)8)13)  
 . Elidan 5) chloram-  
 bucil, steroid cyclosporine  
 chlorambucil  
 1 cyclosporine  
 25%  
 . Cyclosporine  
 5)  
 12)  
 가

REFERENCES

- 1) Kozin F, Houghton V, Bernhard GC. *Neuro-Behçet disease: Two cases and neuroradiologic findings. Neurology* 1977; 27:1148-52.
- 2) Igarashi Y, Watanabe Y, Aso S. *A case of Behçet's disease with otologic symptoms. ORL* 1994;56:295-8.
- 3) Brama I, Fainaru M. *Inner ear involvement in Behçet's disease. Arch Otolaryngol* 1980; 106:215-7.
- 4) Emura A, Takeuchi A, Hashimoto T. *A case of Behçet's disease with Weber's syndrome. J Rheumatol* 1986; 13:459-61.
- 5) Elidan J, Levi H, Cohen E. *Effect of cyclosporine A on the hearing loss in Behçet's disease. Ann Otol Rhinol Laryngol* 1991; 100:464-8.
- 6) Gemignani G, Berrettini S, Bruschini P. *Hearing and vestibular disturbances in Behçet's syndrome. Ann Otol Rhinol Laryngol* 1991; 100:459-63.
- 7) Inaba G. *Clinical features of neuro-Behçet's syndrome. In: Lehner T, Barnes CG, eds. Recent advances in Behçet's disease. London, England: Royal Society of Medicine Services* 1986;235-46.
- 8) Tsunoda I, Kanno H, Watanabe M. *Acute simultaneous bilateral vestibulocochlear impairment in Neuro-Behçet's disease: A case report Auris Nasus Larynx* 1994; 21:243-7.
- 9) Fadli ME, Youssef MM. *Neuro-Behçet's syndrome in the United Arab Republic. Eur Neurol* 1973; 9:76-89.
- 10) Okamoto M, Shitara T, Nishihata S. *Sensorineural hearing loss in Behçet's disease. Nippon Jibiinkoka Gakkai Kaiho* 1982; 85:667-73.
- 11) Alajouanine T, Castaigne P, Lhermitte F. *La meningo-encephalite de la maladie de Behçet. Press Med* 1961; 69:2579-82.
- 12) Pollak L, Luxon LM, Haskard DO. *Labyrinthine involvement in Behçet's syndrome. J Laryngol Otol* 201; 115:522-9.
- 13) Sakurai N, Koike Y, Kaneoke Y. *Sleep apnea and palatal myoclonus in a patient with neuro-Behçet's syndrome. Intern Med* 1993; 32:336-9.